

Original Article

Sero prevalence of Dengue NS-1 Antigen in Tertiary care hospital, Ahmedabad

*Lata R. Patel

Department of Microbiology, Smt. N.H.L. Municipal Medical College, Elishbridge, Ahmedabad - 6

*Correspondence : Email id : drlatapatel76@gmail.com

ABSTRACT

Introduction: Dengue is one of the most serious mosquito-borne viral infections affecting tropical and subtropical countries in the world. Since there is no immune prophylactic or specific antiviral therapy available, timely and rapid diagnosis plays a vital role in patient management and implementation of control measures. The present study was planned to diagnose the dengue infection by detecting dengue NS-1 antigen & to study the seroprevalence of dengue NS-1 antigen

Material and Methods: Dengue NS-1 testing by immunochromatography was performed during september 2011 to november 2012 and the data were analyzed retrospectively. A total 1025 serum samples sent from V.S.G.H. (O.P.D. & Indoor) for the detection of Dengue NS-1Ag.

Results: Total samples tested were 1025 out of which NS-1 seropositive were 167(16.3%). All positive were confirmed by NS 1 ELISA test. Male:female ratio was 2:1. More nos of cases were seen in age group 16- 30 years that is 80(47.9%). Urban:Rural ratio was 4:1. Fever was the commonest presentation in all suspected patients 1025(100%) associated with headache in 935, associated with muscle pain in 906 then fever with headache with muscle pain in 929. Fever with rash in 50, fever with retroorbital pain in 10 and fever with haemorrhagic manifestation in 10 patients were observed. Patients with platelet count less than 50,000 were 33(20%), 50,000 to < 1,00,000 were 60(36%) and > 1,00,000 were 74(44%). According to day of fever, highest nos of seropositive pts were seen in 4th day that is 102(61%) that followed by 3rd day that is 28(16.7%) and from that more nos of seropositive male 61 (36.5%) and seropositive female 41(24.5 %) were seen in 4th day of fever that is followed by on 3rd day 22(13.2%) seropositive male and 6(3.6 %) seropositive female.

Conclusion: New dengue virus strains and serotypes will likely continue to be introduced into urban areas where the densities of *Aedes aegypti* are at high levels. So, for the early and rapid diagnosis NS 1 immunochromatography are very helpful in dengue infection.

Keywords: Dengue infection, NS - 1 protein, rapid diagnostic test

INTRODUCTION: Dengue fever is an important mosquito – borne viral disease of humans. This has been a recurrent phenomenon throughout the tropics in the past decade. Annually, there are an estimated 100 million dengue virus infections worldwide¹. Increasingly, cases of the more severe and potentially lethal dengue hemorrhagic fever (DHF) and dengue shock syndrome (DSS) are reported with children bearing much of the disease burden². The mortality rate of DHF in most countries is 5%, primarily among young children and adults³. Dengue virus is an

enveloped positive sense RNA virus. The genomic RNA is approximately 11 Kb in length and is composed of three structural protein genes that encode for nucleocapsid or core protein(C), a membrane – associated protein (M) , an envelope protein (E), and seven non structural (NS) protein genes including NS 1 protein⁴. Among the non-structural proteins, NS 1 is highly conserved glycoprotein which appears essential for virus replication; no precise function has yet been assigned to it. During acute dengue virus infection, NS 1 is found associated with intracellular organelles or is

transported through the cellular secretory pathway to the cell surface⁵⁻⁷. Now days, detection of NS-1 Ag on rapid tests offer an even faster route to a presumptive dengue diagnosis. NS-1 (Non structural protein) is a highly conserved glycoprotein that is essential for the viability of dengue virus & is produced both in membrane associated & secretory forms by the virus. The detection of secretory NS-1 protein represents a new approach to the diagnosis of dengue infection⁸. It possesses not only group – specific but also type specific determinants and has been recognized as an important antigen in dengue infection⁹.

There are four serotypes of dengue designated dengue 1 through dengue 4 (DEN 1 to DEN 4) that is antigenically related (Monath and Heinz, 1990). Recovery from infection by one serotype can confer life-long protection against that serotype; however, it provides only partial, transient immunity against subsequent infection by the other three dengue serotypes. All four serotypes cause disease, usually asymptomatic or mild dengue fever (Halstead, 1988). Progression from DF to DHF and DSS usually occurs after a second infection with a different serotype, which is due to immune-mediated enhancement of infection, known as antibody-dependent enhancement (ADE) (Halstead, 1988)¹⁰. The present study aim to diagnose the dengue infection by detecting dengue NS-1 antigen, also to study the seroprevalence of dengue NS-1 antigen and to establish their role for early diagnosis.

MATERIAL & METHODS:

Inclusion Criteria:

Clinically suspected Patients experiencing febrile illness consistent with dengue fever with two or more of the following manifestations:

- Headache
- Muscle pain
- Haemorrhagic manifestation

- Retro-orbital pain
- rash

Patients & study design:

Human blood samples from clinically suspected patients of dengue infection were collected in sufficient quantity from september 2011 to Oct. 2012 of the OPD & indoor patients of V. S. General Hospital. A total 1025 serum samples were tested for the detection of NS 1 Ag.

The dengue NS 1 antigen rapid test is an in vitro immunochromatographic, one-step assay designed for the qualitative determination of dengue virus NS 1 antigen in human serum or plasma for the diagnosis early acute dengue infection. The test device contains a membrane strip, which is pre coated with anti dengue NS 1 Ag capture on the test band region. The anti dengue NS 1Ag – colloid gold conjugate and serum move along the membrane chromatographically to the test region “T” and forms a visible line as the antibody-antigen-antibody gold particle complex forms. This test also can detect all 4 dengue serotypes by using a mixture of recombinant dengue envelope proteins.

Remove the test device from the foil pouch and place it on a flat surface. With a disposable dropper, add 3 drops (100µl) of patient’s serum was added into the sample well marked “S” and the test result was interpreted in 15-20 min¹¹.

Interpretation of the SD Bioline Dengue duo rapid test:

The presence of only one colour line within the result window indicated negative result and the presence of two colour lines (“T” band and “C” line) indicated a positive result. When no control line (C) was found the test was considered as invalid¹¹.

OBSERVATIONS & RESULTS: Total 1025 human samples were subjected to Dengue NS 1 Ag by immunochromatography based rapid testing. 167(16.3%) were seropositive for dengue NS 1 Ag. All seropositive samples were confirmed by dengue NS 1 ELISA test. Highest no. of seropositive cases

were noted in September and October and november months of the year 2011 and 2012 (Graph – 1). We observed maximum number of patients coming from urban region 820(80%) of Ahmedabad while 205(20%) were from rural area, so, urban: rural ratio is 4:1 (Graph: 4). More seropositive cases were noted in urban region. Male: female ratio of 1.9: 1 (graph : 3) with maximum seropositive cases were noted in age group 16-30 yrs that is 80(47.9%) shown in Graph - 2.


The most common symptoms apart from fever were headache, muscle pain and rashes and less common retroorbital pain & haemorrhagic manifestations were

observed (Graph: 6). Platelet count was observed < 50,000 in 33(20%), between 50,000 & < 1 lac in 60 (36%) and above 1 lac in 74(44%) shown in Graph - 5.


Duration of fever in days were noted maximum nos of seropositives on 4th day of infection that is 102(61%) followed by 3rd day that is 28(16.7%) after that on later days few no of seropositives were noted (Graph-7).

Days of fever in dengue seropositive male were highest on 4th 61(36.5) and on 3rd day 22(13.2%) while dengue seropositive female were highest on 4th 41(24.5%) and on 3rd 6(3.6%) shown on graph – 8.


Graph – 1 Seasonal variation of dengue infection




Graph – 2 Agewise distribution in dengue suspected & seropositive cases




Graph – 3 Sexwise distribution of dengue infection in suspected & seropositives




Graph – 4 Geographical distribution in dengue suspected & seropositives




Graph – 5 Dengue symptoms in suspected & seropositive cases




Graph – 6 Platelet count in dengue seropositive cases



Graph – 7 Correlation of days of fever with dengue suspected & seropositive cases



Graph 8 : Correlation of days of fever with seropositives gender



DISCUSSION: Dengue infection presents with nonspecific fever that mimics other viral illnesses. The availability of commercial ELISA assays to detect the DEN virus NS 1 protein in acute plasma provides an additional dengue diagnostic tool to the existing approaches of PCR , antibody capture ELISA and less frequently virus isolation ^{7,8,9} . The assessment of NS 1 antigen detection assays as diagnostic tool to the existing dengue diagnostic algorithms. In order to provide timely information for the management of the patients and early public health control of dengue outbreak it is important to establish the diagnosis of acute dengue virus infection during the first few days after manifestation of clinical symptoms.

The seasonality of transmission of dengue with increased activity in the post – monsoon season was seen in the present study ; in accordance with the reported patterns of dengue transmission. Even in the post epidemic period increased dengue virus activity was seen in post monsoon period september to November months of the year 2011 and 2012. These findings indicate that dengue infections are mostly seen in post – monsoon season hence preventive measures should be in full swing at the very onset of the monsoon. Another explanation can be the heavy rains of monsoon season, which usually start in July, August, resulting in stagnant water that serves as breeding ground for vectors of this virus and lead to increased activity in post monsoon period. That is compared with study of Ekta Gupta, Lalit Dar et al¹² in

2006. The reasons may be due to on these months large stores of water. Also the breeding habit of *Aedes aegypti* is highest during pre and post monsoon period. But sporadic cases extend up to December which indicates endemicity of the infection up to December.

Female aedes mosquito, the vector of the virus is peridomestic in nature. The tropical zones of the world having monsoon rains are the usual habitat of this vector⁹. The breeding of *Aedes aegypti* is highest during pre and postmonsoon period¹⁰. We found the epidemic in post monsoon period. Major contributory factors to this increased activity may be due to changes in weather pattern such as El-Nino phenomenon².

In present study, higher nos of males were affected than females and maximum number of cases were between 16 – 30 years of age group observed that is correlated with study done by Halstead¹⁹ had pointed out as early as 1970 that males predominate among those with milder disease but females account for more severe illness. He suggested that either immune response in females are more competent than in males, resulting in greater production of cytokines, or the capillary bed of females is prone to increased permeability. Kaplan¹⁷ in Mexico suggests that an incidence bias in favour of females is related to the timing of the survey interviews, while Goh²⁰ puts forward that low incidence among women occurs because they stay at home and are less exposed to infection^{17,18}. Gupta *et al*¹² from India showed a maximum number of cases between the ages of 21 to 30 years.

In present study, urban population were highly affected that is 80% than rural population having 20%

.Historically, DF/DHF has been reported as occurring predominantly among urban populations where density of dwellings and short flying distance of the vector create the right conditions for transmission²¹. Increased transport contact, mobility and spread of periurbanisation have been the most frequently cited reasons for spread of dengue to rural areas²².

Data obtained in this study show DEN virus antigens were detected from as early as Day 1 (5.8% of samples) up to Day 9 (0.4% of samples) of fever. These findings are comparable to a study by Alcon *et al*²³ in 2006 who recovered NS1 antigen until day 9 of symptoms. Antigen detection was highest between Days 3 and 4 with a detection rate ranging from 12 % to 38 %.

CONCLUSION: High prevalence rate in our region particularly in pre monsoon and monsoon season gives an alarm to the doctors regarding early and accurate diagnosis of dengue virus infection and its complications. Prompt diagnosis of evaluation of rapid dengue NS1 antigen test showed that this test is highly appropriate for diagnosis of dengue infection as it has proven to be a rapid, easily applicable, sensitive and specific method. NS1 antigen detection has potential value for screening patient samples during the early acute phase. It is rapid, easily be performed, interpreted early and has a extended shelf life. We conclude that rapid test is an effective tool, if when used in combination with NS 1 MAC ELISA in single sample of suspected cases, has the ability to improve the diagnostic algorithm contributing significantly to clinical treatment and to control dengue viral infection.

REFERENCES:

1. Gubler DJ, Metzger M. Impact of dengue/dengue haemorrhagic fever on the developing world. *Adv Virus Res* 1999; 53:35-70.
2. Whitehead SS, Blaney JE, Durbin AP, Murphy BR. Prospects for dengue virus vaccine. *Nat Rev Microbiol* 2007;5:518-28

3. World Health Organisation. Key Issues in dengue vector control towards the operationalisation of a global strategy: report of consultation . Geneva : WHO;1995.(CTD/FIL(DEN)/IC.96.1)
4. Clyde K, Kyle JL,Harris E. Recent advances in deciphering viral and host determinants of dengue virus replication and pathogenesis. *J Virol* 2006;80: 11418-31.
5. Mackenzie JM,Jones MK,Young PR. Immunolocalisation of the dengue virus non-structural glycoprotein NS 1 suggests a role in viral replication. *Virology* 1996: 220: 232-40
6. Guzman MG, Kouri G. Advances in dengue diagnosis. *Clinn Diagn Lab Immunol* 1996;3:621-7.
7. Blacksell SD,Mammom MP Jr,Thongpaseutha S,Gibbons RV, Jarman RG et al. Evaluation of the panbio dengue virus non-structural antigen detection and IgM antibody enzyme linked immunosorbent assaya for the diagnosis of acute dengue infections in Laos. *Diagn Microbiol Infect Dis* 2008;60 :43-9.
8. Dussart P, Labeau B , Lagathu G, Louis P, Nunes MRT, Rodrigues SG, et al. Evaluation of an Enzyme Immunoassay for detection of dengue virus NS 1 antigen in human serum. *Clin Vaccine Immunol* 2006; 13:1185-
9. Kumarasamy V, Wahab AH,Chua SK, Hassan Z, Chem YK, Mohmad M, et al. Evaluation of a commercial dengue NS 1 antigen capture ELISA for laboratory diagnosis of acute dengue virus infection. *J Virol Methods* 2007,140:75-9.
10. Sekaran , S.D.,Ew, C.L., Subramaniam, G., Kanthesh , B.M.Sensitivity of dengue virus NS 1 detection in primary and secondary infections . *African J of Microbiology research* vol. 3(3),pp 105-110 march 2009
11. STANDARD DIAGNOSTICS , INC. SD BIOLINE dengue Duo Dengue NS 1 Ag + Ab combo
12. Ekta Gupta, Lalit Dar, Geetanjali Kapoor & Shoba Broor. The changing epidemiology of dengue in Delhi, India . *Virology Journal* 2006, 3:92
13. World Health organization. Prevention & control of dengue & Dengue haemorrhagic fever . Disease SEARO publication No. 29, New Delhi 1999
14. Yunus EB National guidelines for clinical management of dengue & dengue haemorrhagic fever . Disease control directorate DGHS, Dhaka-2000
15. World Health organization. Dengue in the Who western pacific region wkly , *Epid Rec.* 1998, 73 : 273-280
16. Shekar KC , Huat Ol: Epidemiology of Dengue/dengue haemorrhagic fever in Malaysia- a retrospective epidemiological study 1973-1987. Part 1 : *DHF Asia Pac J Public Health* 1992, 6: 15-25
17. Kaplan JE,Eliason DA, Moore M Epidemiological investigations of dengue infection in mexico 1980. *Am J Epidemiol* 1983,117: 335-343
18. Kabra SK ,Jain Y .Pandey Rm Dengue haemorrhagic fever in children in the 1996 delhiepidemic. *Trans R Soc Trop Med Hyg* 1999,93 : 294-298
19. Halstead SB, Nimmannitya S, observations related to pathogenesis of DHF. IV. Relation of disease severity to antibody response & virus recovered. *Yale J Biol Med* 1970, 42: 311-328
20. Goh KT : Changing epidemiology of dengue in Singapore. *Lancet* 1995, 346: 1098.
21. Muto RSA : Df/DHF & its control status in WHO's western pacific region by 1999. In WHO internal report Manilla , WHO western Pacific regional Office, 2004:4
22. Githeko AK ,Lindsay SW : Climatic change & vector -borne disease : a regional analysis . *Bull World Health Organ* 2000,78: 1136-1147

23. Alcon S, Talarmin A, Debruyne M, Falconar A. Enzyme linked immunosorbent assay specific to dengue virus type 1 Nonstructural protein NS1 reveals circulation of the antigen in the blood during acute phase of disease in patients experiencing primary or secondary infections. J Clin microbial 2002: 40:376-81

Date of submission: 22 February 2013
Date of Provisional acceptance: 17 March 2013
Date of Final acceptance: 28 April 2013
Date of Publication: 03 June 2013
Source of Support: Nil ; Conflict of Interest: Nil

International Journal of Healthcare & Biomedical Research

Official Publication of **IJBAMR FORUM**

..... *Ahead of prints*

Salient features:

- Regular publishing in the first week of January , April, July & October
- Wide Indexing & Abstracting
- Wide readership across the world
- Available in electronic version
- Open access Biomedical Journal
- Officially listed in HIFA 2015
- Officially listed in IJCME
- Listed in Google scholar
- Proposal send for HONCode & Medhunt accreditation
- Proposal send for IC Value
- Rapid publication process
- Add on facility available

Official web portal: www.ijhbr.com