

A STUDY ON BACTERIAL HAEMORRHAGIC SEPTICEMIA IN FARMED YOUNG RUSSIAN STURGEON IN TURKEY (*Acipenser gueldenstaedtii*)

Gulsen TIMUR^{*}; Tulay AKAYLI[†]; Jale KORUN[†]; R. Eda YARDIMCI^{*}

Makalenin alındığı tarih: 04, 2009
Kabul tarihi: 12, 2009

ABSTRACT

Bacterial haemorrhagic septicemia outbreaks occurred in young farmed Russian sturgeon (*Acipenser gueldenstaedtii*) in 2002-2003 associated with *Aeromonas hydrophila* alone or mixed infection with *Flavobacterium hydatis*. These bacteria induced low mortalities among the fish reared in concrete ponds at Sapanca Freshwater Fish Culture and Research Station, Istanbul University, Fisheries Faculty. Clinically, affected fish, 1-2 years old (100-380 g in weight) showed haemorrhages on the skin of ventral side of head, at the base of fins and around ganoid scales. Affected fish gills generally were congested, haemorrhagic and necrotic. Some affected fish gill filaments showed epithelial cell hyperplasia, necrosis and sloughed of the necrotics cells.

Key Words: *Aeromonas hydrophila*, *Flavobacterium hydatis*, Russian sturgeon, haemorrhagic septicemia.

ÖZET

TÜRKİYE'DE GENÇ KÜLTÜR RUS MERSİN BALIKLARINDA (*Acipenser gueldenstaedtii*) GÖRÜLEN BAKTERİYEL HEMORAJİK SEPTİSEMİ ÜZERİNDE BİR ÇALIŞMA

Istanbul Üniversitesi Su Ürünleri Fakültesi Sapanca İçsu Balıkları Üretim ve Araştırma İstasyonu'nda beton havuzlarda yetiştiriciliği yapılan genç Rus mersin balıklarında (*Acipenser gueldenstaedtii*) 2002-2003 yıllarında *Aeromonas hydrophila* tek başına veya *Flavobacterium hydatis* ile karma enfeksiyon şeklinde bakteriyel hemorajik septicemia oluşturarak balıklarda düşük mortaliteye neden olmuştur. Klinik olarak hasta balıklarda (100-380gr) deri renginde koyulaşma, yüzgeç diplerinde ve ganoid pulların çevresinde hemorajiler görülmüştür. İç organlarda hiperemi ve hemoraji yaygın bir şekilde görülürken solungaçların genellikle hiperemik, hemorajik ve nekrotik olduğu tespit

edilmiştir. Bazı hasta balıkların solungaç filamentlerinde epitelyal hücre hiperplazisi, nekroz ve nekrotik hücrelerde dökülme görülmüştür.

Anahtar Kelimeler: *Aeromonas hydrophila*, *Flavobacterium hydatis*, Rus mersini, hemorajik septicemia.

INTRODUCTION

Despite the extensive information about the bacterial disease in most teleost fishes, a few data exist on bacterial diseases of sturgeon. A motile aeromonad septicemia caused by *Aeromonas hydrophila* frequently induced considerable losses in Persian sturgeon (*Acipenser persicus*) fingerlings in northern Iran (Soltani and Kalbassi, 2001) and also in the Harrison river sturgeon in Canada (Raverty and Nikl, 1999). *Yersinia ruckeri* was found responsible for 10% mortalities in the young (15-30 g) cultured sturgeon (*Acipenser baeri*) in south western of France (Vuillaume *et al.*, 1987). Bauer *et al.*, (2002) reported that *Flavobacterium johnsoniae* was isolated from diseased farmed young sturgeons (3-4 g) in Russia and it was also isolated from 6-7 years old diseased farmed Russian sturgeon with low mortality in Turkey reported by Karataş *et al.*, 2010. An outbreak of *Pseudomonas fluorescens* was reported in young (10g in size) farmed Siberian sturgeon (*A. baeri*) with high mortality (40%) in northern Italy (Brunetti *et al.*, 2006).

The aim of this study was to characterize pathogens responsible for the outbreaks of disease in young farmed Russian sturgeon (1-2 years old) reared from fertilized eggs which were imported from Krasnodar Russia in 2001. The result of bacteriological examinations and details of the histopathology of the disease are presented here.

MATERIAL AND METHODS

Fish

Ten affected fish (100-380g) obtained from concrete ponds of Istanbul University, Fisheries Faculty Research Station in Sapanca in 2002-2003, was sampled for bacteriology and histopathology. Water temperature at the time of sampling varied from between 20 and 22°C.

Bacteriology

Bacteriological samples of liver, kidney and spleen were streaked onto Tryptic Soy Agar (TSA) plates and plates were incubated at 22°C for 48h. The morphological and physiological characteristics of bacterial colonies from each plates were determined by conventional biochemical and physiological tests (Austin and Austin, 1987) and bacteria identified following Bergey's Manual (Holt *et al.*, 1994).

Histopathology

Samples of tissues from gut, kidney, spleen, liver heart and gills were processed for histopathology by fixing in 10% buffered formalin, and processed for paraffin embedding. Histological sections (5µm) were stained with haematoxyline and eosin and examined by light microscopy (Bullock, 1978).

^{*} Istanbul University, Faculty of Fisheries, Department of Aquaculture, 34470, Laleli -Istanbul, Turkey.

^{*} Corresponding author's E-mail: gulsentimur@yahoo.com

[†] Akdeniz University, Faculty of Fisheries, Department of Aquaculture, Campus, 07049, Antalya, Turkey.

RESULTS

Affected fish (100-380g) generally showed darkening of the skin, with haemorrhages on the ventral side of the head skin, at the base of fins and around the ganoid scales (Figure 1). Considerable level of mortalities occurred in fish stocks.



Figure 1. Hyperemia and haemorrhages on the ventral side of the head skin, at the base of fins and around the ganoid scales in the affected fish.

Şekil 1. Hasta balıklarda başın alt yüzünde, yüzgeç diplerinde ve ganoid pulların etrafında hiperemi ve hemorajı

Internally, the most obvious findings were congestion of the visceral organs with haemorrhages over the viscera (Figure 2). In addition to these findings, black colouration of the liver was also observed in some affected fish.

Two Gram-negative bacteria cultured from visceral organs following incubation of kidney, spleen and liver swabs at 22°C for 48h on TSA. These bacteria were identified as *A. hydrophila* and *Flavobacterium hydatis*. *A. hydrophila* produced cream coloured colonies consist of Gram negative, motile and short (0.5-0.8 x 1.5-3.5µm) rods, were not sensitive vibriostat (O/129); cytochrome oxidase; and catalase positive, fermentative by O/F glucose test. *F. hydatis* produced yellow coloured colonies consist of long rods (0.5x 8-10µm), showed gliding movements, catalase positive, but not oxidase positive, fermentative by O/F glucose test. A few long, slender thread-like cells were seen in the young cultures of the *F. hydatis* in Gram stained preparates (Figure 3). The morphological, physiological and biochemical features of the isolates were shown in Table 1.



Figure 2. Congested visceral organs, with haemorrhages over the visera.

Şekil 2. Hiperemik visceral organlar ile visceral organlarda hemorajiler




Figure 3. A few long, slender thread-like cells (arrowed) in young cultures of *Flavobacterium hydatis* (Gram x1000).

Şekil 3. *Flavobacterium hydatis*'in genç kültürlerinde bir kaç ince uzun iplik benzeri (okla gösterilmiştir) bakteri hücreleri (Gram x1000).

Table 1. The morphological, physiological and biochemical features of isolated bacteria.
Tablo 1. İzole edilen bakterilerin morfolojik, fizyolojik ve biyokimyasal özellikleri

	<i>Aeromonas hydrophila</i>	<i>Flavobacterium hydatis</i>
Gram staining	-	-
Motility	+	+*
Cytochrome Oxidase	+	-
Oxidative-Fermentative Metabolism	+/+	+/+
Voges Proskauer reaction	-	V
Methyle red test	V	-
Indole	V	V
Citrate	+	-
Arabinose	-	V
Galactose	-	+
Mannose	+	+
Lactose	-	-
Fructose	+	-
Maltose	+	+
Sucrose	+	+
Glucose	+	+
Mannitol	-	-
Xylose	-	V
Arginine dihydrolase	+	+
Lysine decarboxylase	-	-
Ornithine decarboxylase	-	-
Growth 37°C	+	+
%0 (w/v) NaCl	+	+
%4 (w/v) NaCl	-	-
%7 (w/v) NaCl	-	-
O/129	R	S
ONPG (β-Galaktosidase)	+	+

*: gliding motility; V: variable; R: resistance; S: sensitive

Histologically, diseased **sturgeons** showed depletion of haemopoietic elements both in the renal (Fig. 4a) and splenic haemopoietic tissue. The remaining cells of spleen were usually necrotic. The kidneys showed peritubular and glomerular oedema, considerable tubular degeneration and necrosis (Figs 4b). Multifocal large liquefactive necrotic areas and haemorrhages were found in the affected liver and focal necrosis is found in the cardiac muscle. Multifocal melanomacrophage centers and liquefactive necrotic areas were seen in some affected liver (Fig. 4c) sections. The intestinal (Fig. 4d) and gastric mucous membrane was usually necrotic and sloughed into the lumen. Affected fish gills with *A. hydrophila* were generally congested and haemorrhagic (Fig. 4e). Epithelial hyperplasia, sloughed epithelial cells and mucus were observed in **the** gill filaments of affected fish with *F. hydatis* (Fig. 4f).




Figure 4. Histological sections of affected fish tissues (a) depletion of haemopoietic elements, degenerative and necrotic tubules in kidney (H&E, X100), (b) periglomerular oedema and haemorrhage in the affected glomerulus (H&E, X200), (c) multifocal melanomacrophage centers in the affected liver (H&E, x100), (d) necrotic mucosa membrane and sloughed into the lumen of the affected intestine (H&E, x200), (e) congested and haemorrhagic gill filaments (H&E, x200), (f) and epithelial hyperplasia, haemorrhages of the necrotic gill filaments tips (H&E, x100).

Şekil 4. Hasta balık dokularının histolojik kesitleri (a) böbrekte hemopoietik dokuda azalma, dejeneratif ve nekrotik tübüller (H&E, X100), (b) glomeruluslarda periglomerular ödem ve hemoraji (H&E, X200), (c) karaciğerde multifokal melanomakrofaj odakları (H&E, x100), (d) bağırsakta nekrotik mukoza membranında lümene dökülme (H&E, x200), (e) solungaç filamentlerinde hiperemi ve hemoraji (H&E, x200), (f) nekrotik solungaç filament uçlarında epitelyal hiperplazi ve hemorajiler (H&E, x100).

DISCUSSION

For the most diseases investigation of sturgeon, regardless of species, a bacterial agent has been identified as the definitive cause of mortality in many countries (Vuillaume *et al.*, 1987; Raverty and Nikl, 1999; Francis-Floyd, 2000; Soltani and Kalbassi, 2001; Brunetti *et al.*, 2006).

Francis-Floyd (2000) has been reported that most bacterial agents isolated from most sturgeon disease cases which were predisposed stress factor in their history, were examined University of Florida; identified as opportunistic pathogen bacteria namely, *A. hydrophila*, *Pseudomonas* spp. and *A. sobria*. In this study, similarly we also identified opportunistic pathogen bacteria namely *A. hydrophila* and *F. hydatis* as definitive causes of mortality of Russian Sturgeon reared in the large concrete ponds at low level of dissolved oxygen (<4mg/l) with organic loads, particularly when the accumulations are on the bottom of ponds.

The affected fish had showed darkening in colour with haemorrhages on the ventral side of the head skin, and at the base of fins and around the large ganoid scales and congested visceral organs, with haemorrhages over the viscera, mortality, as described in other teleost fish infected with *A. hydrophila* (Richards and Roberts, 1978, Soltani and Kalbassi, 2001; Guvener and Timur, 2005).

F. hydatis isolates were consist of 8-10µm long rods as described other workers (Strohl and Tait, 1978; Austin and Austin, 1987, Bernardet *et al.*, 1996) but we observed a few long, slender thread-like cells in young cultures. *F. hydatis* exhibits swarming and gliding motility under certain conditions, and the structure of its cell wall is similar to that of *F. johnsoniae* (Bernardet *et al.*, 1996, Karataş *et al.*, 2010). Their micromorphology, gliding motility and the ability of the acid production from glucose, cytochrome oxidase negative and fermentative (F) metabolism character had made them distinct from other *Flavobacterium* spp. (Bernardet *et al.*, 1996, Austin and Austin, 1999). For the reason *F. johnsoniae* should have given positive cytochrome oxidase reaction we differentiated our isolate as *F. hydatis*.

Depletion of haemopoietic elements in the renal and splenic haemopoietic tissue, considerable tubular degeneration and necrosis in the kidney, multifocal necrotic areas and haemorrhages in the liver, in the gills and necrotic intestinal and gastric mucous membrane were also similar histopathological findings were described in other teleost fish with motile aeromonad septicemia (Richards and Roberts, 1978; Guvener and Timur, 2005). In addition to these findings in the affected fish with *F. hydatis*; hyperplasia of the gill epithelia, necrose and sloughed of epithelial cells were observed beside the haemorrhagic as described in the bacterial gill diseases induced by *flavobacteria* (Noga, 1999). Diffuse melanomacrophage centers were observed in some affected liver sections similar to the kidney which that normally do not observe in the teleost fish liver. **These diffuse** melanomacrophage centers should be turned the liver colour to black.

Finally, this is the first report of the haemorrhagic septicaemia induced by *A. hydrophila* associated with or **without** *F. hydatis*, as a secondary pathogen in farmed Russian sturgeon in Turkey.

REFERENCES

- Austin B and Austin DA (1999). "**Bacterial Fish Pathogen: Disease in Farmed and Wild Fish.**" Ellis Harwood Limited, England, pp 63-80, 91-98. ISBN 085312844 8.
- Bauer ON, Pugachev ON and Voronin VN (2002). Study of parasites and disease of sturgeons in Russia: a review. *J. Appl. Ichthyol.* **18**, 420-429.
- Bernardet JF, Segers P, Vancanneyt M, Berthe F, Kersters K and Vandamme P (1996). Cutting Gordian Knot: Emended Classification and Description of the Genus *Flavobacterium*, Emended Description of the Family *Flavobacteriaceae*, and Proposal of *Flavobacterium hydatis* nom. nov. (Basonym, *Cytophaga aquatilis* Strohl and Tail 1978). *International Journal of Systematic Bacteriology*, **46** (1), 128-148.
- Brunetti R, Gasparri F, Marturano S and Prearo M (2006). *Pseudomonas fluorescens* infection in farmed Siberian sturgeon (*Acipenser baeri*). *Ittiopatologia*, **3**, 221-226.
- Bullock AM (1978). "Laboratory Methods. **"In Fish Pathology"**. Edited. By Ronald J. Roberts. Bailliere Tindall, London, pp 235-275 .
- Francis-Floyd R (2000). Diseases history of cultured sturgeon in Florida, 1990-1999, Proceedings of the Florida sturgeon culture risk assessment workshop. pp : 33-37.
- Guvener RP and Timur G (2005). A study on determination of the Aeromonad infections in some aquarium fish. *Istanbul University Journal of Fisheries and Aquatic Sciences*, **19**, 27-39.
- Holt J, Krieg NR, Sneath RAH, Staley JI, Williams ST (1994). "**Bergey's Manual of Determinative Bacteriology"**. Ninth edition Williams and Wilkins. ISBN 0 683 006037.
- Karatas S, D. Ercan, T. M. Steinum, E. Turgay, D. Memis and A. Candan (2010). First isolation of a *Flavobacterium johnsoniae* like bacteria from cultured Russian sturgeon in Turkey. *Journal of Animal and Veterinary Advances*, **9** (14), 1943-1946.
- Noga EJ (1999). "**Fish Disease Diagnosis and Treatment"**. Iowa State University Press. Blackwell Publishing Company, Iowa. ISBN 0-8138-2558-X.
- Raverty S and Nikl L (1999). Sturgeon mortalities along the Harrison river, British Columbia. Animal health center, Newsletter, *Diagnosis and Diary*, **9**, 10-11.
- Richards RH and Roberts RJ (1978). The bacteriology of Teleost. "**In Fish Pathology"**. Edited. By Ronald J. Roberts. Bailliere Tindall, London, pp 183-204.
- Soltani M and Kalbassi MR (2001). Protection of Persian surgeon (*Acipenser persicus*) fingerling against *Aeromonas hydrophila* septicemia using three different antigens. *Bull. Eur. Ass. Fish Pathol.* **21** (6), 235-240.
- Strohl WR and Tait LR (1978) *Cytophaga aquatilis* sp. nov., a Facultative anaerobe isolated from the Gills of Freshwater Fish. *Int. J. Syst. Bacterial*, **28**, 293-303.
- Vuillaume A, Brun R, Cnene P and Lesel R (1987). First Isolation of *Yersinia ruckeri* From Sturgeon, *Acipenser baeri* BRANT, in South West of France. *Bull. Eur. Ass. Fish Pathol.* **7** (1), 19.